


Duration: 3 hours

Marks: 75

I. Do as directed**20 M**

1. Predict m/z of molecular ion of ethyl bromide
2. Predict m/z of base peak for acetaldehyde
3. Predict the number of signals in the ^{13}C NMR of cyanobenzene
4. Give an example of a molecule containing D_2O exchangeable proton/s
5. Calculate index of hydrogen deficiency for $\text{C}_7\text{H}_7\text{NO}_2$
6. Predict the number of signals in ^1H NMR of



7. Predict the characteristic IR frequencies of -



8. Will this compound undergo McLafferty rearrangement?



9. Give one application of GC AAS spectroscopy
10. Give examples of any two sorbents used in HPTLC plates
11. Name any two anionic exchangers
12. Give any one difference between Flash chromatography and HPLC.
13. Name the mobile phase used in supercritical fluid chromatography
14. Define 'J' with respect to NMR spectroscopy.
15. What is plotted on X-axis and Y-axis in HETCOR spectrum?
16. What is Stokes scattering with respect to Raman spectra?
17. Which radioactive material is used in RIA of digoxin?
18. Draw DSC curve with proper nomenclatures.
19. What increment will you add to the parent value for the exocyclic double bond marked in the structure?



20. Arrange the following in increasing order of frequency for $\text{C}=\text{O}$ stretch
Aliphatic ketones, α,β -unsaturated ketone, $\alpha\beta$ -unsaturated amide

II. Long answer questions- (Answer any two)

20M

1. a. A compound with molecular weight 104 has the following spectral characteristics:

6M


IR – 3125-2857, 1718, 1449 cm^{-1}

$^1\text{H NMR}$ – δ 10.95 (bs, 5.4sq), 4.13(s, 11.0sq), 3.66(q, $J=7.1\text{Hz}$, 10.6sq),

1.27 (t, $J=7.1\text{Hz}$, 16.2sq). Predict structure. Justify your answer.

- 1.b. How will you distinguish between the following compounds using spectral techniques?

4M



2. a. Elucidate the structure of compound with molecular formula $\text{C}_{10}\text{H}_{11}\text{O}_2\text{Cl}$

6M

and the following spectral details:

IR - 1745, 1600, 1580 cm^{-1}


$^1\text{H NMR}$ – δ 2.00(s, 3H), 2.8(t, 2H, $J=6\text{Hz}$), 4.1(t, 2H, $J=6\text{Hz}$), 7.1(d, 2H, $J=8\text{Hz}$),

7.3(2h, d, $J=8\text{Hz}$).

Deduce structure and justify your answer.

2. b. Predict the mass spectrum of the following compound depicting at least two fragmentation pathways-one depicting fission and another involving rearrangement.

4M



3. a. Deduce the structures from the given $^{13}\text{C NMR}$ spectra-

6M



(i) M. Wt-88 13.8(q), 20.0(q), 60.0(t), 170.3(s)

(ii) M. Wt. 100 30.2(q), 52.8(t), 201.9(s)

(iii) Mol. Formula- $\text{C}_{10}\text{H}_{12}\text{O}$ 13.7(q), 17.6(t), 40.2(t), 127.9(d), 128.8(d), 133(d), 137(d), 199(s)

3.b. Explain how you will distinguish between-


4M



III. Short answer questions- (Answer any seven)

35M

1. Predict the λ_{max} in the UV spectra of both the structures given below-



2. Explain NOESY and COSY techniques in detail

3. Predict IR values-



4. Explain the importance of LC-MS in brief and describe one interface

5. Explain the terms in brief- a) Metastable ions (b) Isotopic ions

6. Elaborate on ELISA technique

7. What is meant by supercritical fluid chromatography. Give 2 advantages of the technique over HPLC.

8. Write principle and instrumentation for TGA

9. Explain ion chromatography in brief

CHARACTERISTIC INFRARED ABSORPTIONS OF FUNCTIONAL GROUPS

GROUP	RANGE	Intensity	Range cm ⁻¹
Unsaturated vinyl ester type	3.36-5.65	(s)	1800-1770
α,β -unsaturated aldehyde	5.78-5.82	(s)	1730-1717
α,β -unsaturated ketone	5.78-5.82	(s)	1730-1717
β,γ -unsaturated ketone	5.68-5.75	(s)	1700-1740
α,β -unsaturated ketone	5.56	(s)	1800
α,β -ketone (enol)	5.70-5.75	(s)	1735-1740
Carbonate	5.06	(s)	1850
5. CARBOXYLIC ACIDS			
a. Carbonyl stretching vibrations saturated aliphatic	5.20-5.28	(s)	1715-1700
α,β -unsaturated aliphatic	5.83-5.92	(s)	1715-1690
aryl	5.82-5.95	(s)	1700-1680
b. Hydroxyl stretching (bonded)			
several bands	3.70-4.00	(w)	2700-2500
	6.21-6.45	(s)	1610-1550
	and 7.15-7.69	(s)	1400-1300
c. Carboxylic anion stretching			
5. ANHYDRIDES/STRETCHING VIBRATIONS			
a. Saturated cyclic	5.41-5.56	(s)	1850-1800
	and 5.95-5.75	(s)	1780-1740
	and 5.47-5.62	(s)	1830-1780
b. α,β -Unsaturated secondary cyclic			
	and 5.65-5.81	(s)	1770-1720
	and 5.35-5.49	(s)	1870-1820
c. Saturated 5-membered ring			
	and 5.56-5.71	(s)	1800-1750
	and 5.41-5.56	(s)	1850-1800
	and 5.47-5.62	(s)	1850-1710
6. ACYL HALIDES/STRETCHING VIBRATIONS			
a. Acyl fluorides	5.41	(s)	1850
	and 5.57	(s)	1795
b. Acyl chlorides	5.53	(s)	1810
c. Acyl bromides	5.61-5.72	(s)	1780-1750
d. α,β -Unsaturated primary	5.72-5.82	(m)	1750-1720
	and 5.19	(s)	1925
	and 5.47	(s)	1825
	and 5.47	(s)	1825
e. COF ₂			
f. COCl ₂			
g. COBr ₂			
7. AMIDES			
a. Carbonyl stretching vibrations			
Primary, solid and concentrated solution	5.95-6.14	(s)	1650-1630
Primary, dilute solution	5.95-6.14	(s)	1650-1630
Secondary, solid and concentrated solution	5.83-5.99	(s)	1700-1670
Secondary, dilute solution	5.99-6.24	(s)	1670-1630
Tertiary, solid and all solutions	5.95	(s)	1680
Cyclic, β -lactams, dilute solution	5.88	(s)	1700
Cyclic, γ -lactams, dilute solution			

TURN OVER

CHARACTERISTIC INFRARED ABSORPTIONS OF FUNCTIONAL GROUPS

GROUP	RANGE	Intensity	Range cm ⁻¹
Cyclic, γ -lactams, fused to another ring, dilute solution	5.71-5.88	(s)	1750-1700
Cyclic, β -lactams, dilute solution	5.68-5.78	(s)	1760-1730
Cyclic, β -lactams, fused to another ring, dilute solution	5.62-5.65	(s)	1780-1770
Ureas, cyclic	6.02	(s)	1660
Ureas, cyclic, 6-membered ring	6.10	(s)	1640
Ureas, cyclic, 5-membered ring	5.81	(s)	1730
Urethane	5.75-5.92	(s)	1700-1690
Imides/acyclic			
	and 5.55	(s)	1710
	and 5.83	(s)	1710
	and 5.85	(s)	1715
Imides, cyclic, 6-membered ring			
	and 5.78	(s)	1720
	and 5.99	(s)	1670
	and 5.65	(s)	1770
	and 5.88	(s)	1720
Imides, cyclic, α,β -unsaturated, 6-membered ring			
	and 5.59	(s)	1790
	and 5.85	(s)	1710
Imides, cyclic, α,β -unsaturated, 5-membered ring			
	and 5.25	(m)	1850
	and 5.24	(m)	1820
	and 5.89	(m)	1810
	and 5.15	(m)	1810
	and 5.22	(m)	1820
	and 5.12	(m)	1850-1810
Secondary, free base band			
	and 6.17-6.29	(s)	1620-1590
	and 6.45-6.62	(s)	1590-1510
Primary amides, dilute solution			
Secondary amides, dilute solution			
Secondary amides, dilute solution			
Miscellaneous chromophoric groups			
1. ALCOHOLS/AND PHENOLS			
a. O-H stretching vibrations			
Free O-H	2.74-2.79	(v,sh)	3600-3500
Intermolecularly hydrogen bonded (change on dilution)			
single bridge compounds	2.82-2.90	(v,sh)	3550-3450
polymers	2.94-3.13	(v,sh)	3400-3200
polymeric association			
Intramolecularly hydrogen bonded (no change on dilution)			
single bridge compounds	2.82-2.90	(v,sh)	3550-3450
chelic compounds	3.1-4.0	(v,sh)	3300-3100

Abbreviations: s = strong, m = medium, w = weak, v = variable, sh = broad, sh = sharp, ~ = approximately

CHARACTERISTIC INFRARED ABSORPTIONS OF FUNCTIONAL GROUPS

Group	Range μ	Intensity	Range cm^{-1}	
b. O-H Bonding and C-O stretching vibrations Primary alcohols	and 9.5	(s)	1050	
	and 7.4-7.9	(s)	1350-1260	
	and 9.1	(s)	1100	
	and 7.4-7.9	(s)	1350-1260	
	and 8.7	(s)	1150	
Tertiary alcohols	and 7.1-7.6	(s)	1410-1310	
	and 8.3	(s)	1200	
Phenols	and 7.1-7.6	(s)	1410-1310	
	and 7.1-7.6	(s)	1410-1310	
2. AMINES				
	a. N-H stretching vibrations Primary; free; two bands	and 2.86	(m)	3500
		and 2.94	(m)	3400
		2.86-3.02	(m)	3500-3310
		2.94-3.03	(m)	3450-3300
		3.2-3.3	(m)	3100-3030
	b. N-H Bending vibrations Primary Secondary Amine salts	6.08-6.29	(s-m)	1650-1590
		6.08-6.45	(w)	1650-1550
		6.25-6.35	(s)	1600-1575
		and 6.67	(s)	1500
7.46-8.00		(s)	1340-1250	
c. C-N Vibrations Aromatic, primary Aromatic, secondary Aromatic, tertiary Aliphatic	7.41-7.81	(s)	1350-1290	
	7.36-7.64	(s)	1360-1310	
	8.2-9.3	(w)	1220-1020	
	and 7.1	(w)	1410	
	and 7.1	(w)	1410	
3. UNSATURATED NITROGEN COMPOUNDS				
	a. C=N stretching vibrations Alkyl nitriles α,β -Unsaturated alkyl nitriles Aryl nitriles Isocyanates Isocyanides	5.72-4.46	(m)	2260-2240
		4.47-4.51	(m)	2235-2215
		4.46-4.50	(m)	2240-2220
		4.40-4.46	(m)	2275-2240
		4.50-4.83	(m)	2220-2070
	b. C=N - stretching vibrations (imines, oximes) Alkyl compounds α,β -Unsaturated compounds C=N=N - stretching vibrations and compounds	5.92-6.10	(v)	1690-1640
		6.02-6.14	(v)	1660-1630
		6.14-6.35	(v)	1620-1575
		4.64-4.70	(s)	2135-2130
4.63-4.72		(s)	2160-2120	
c. N=N stretching vibrations and N=N stretching vibrations dimides	and 7.46-8.48	(w)	1340-1180	
	and 7.46-8.48	(w)	1340-1180	

CHARACTERISTIC INFRARED ABSORPTIONS OF FUNCTIONAL GROUPS

Group	Range μ	Intensity	Range cm^{-1}		
f. C-NO ₂ , Nitro compounds; aromatic	6.37-4.67	(s)	1570-1500		
	and 7.30-7.79	(s)	1370-1300		
	5.37-4.45	(s)	1570-1500		
	and 7.25-7.29	(s)	1300-1170		
	6.06-4.35	(s)	1550-1600		
h. C-NO, Nitroso compounds	and 7.70-8.00	(s)	1300-1250		
	6.25-6.57	(s)	1600-1550		
	5.95-6.08	(s)	1650-1600		
i. O-NO, Nitrites	and 6.15-6.27	(s)	1625-1610		
	and 6.15-6.27	(s)	1625-1610		
4. HALOGEN COMPOUNDS, C-X STRETCHING VIBRATIONS					
	a. C-F	7.1-10.9	(s)	1400-1000	
	b. C-Cl	12.5-16.8	(s)	800-600	
	c. C-Br	16.5-20.0	(s)	600-500	
	d. C-I	~30	(s)	~500	
	5. SULFUR COMPOUNDS				
		a. S-H stretching vibrations	3.3-3.32	(w)	2800-2550
		b. C-S stretching vibrations	4.35-4.52	(s)	2300-1950
		c. S=O stretching vibrations; sulfoxides	9.35-9.71	(s)	1070-1030
		d. S=O stretching vibrations; sulfones	8.62-8.77	(s)	1150-1140
and 7.41-7.69		(s)	1350-1300		
e. S-O stretching vibrations; sulfites		8.13-8.70	(s)	1230-1150	
and 7.00-7.21		(s)	1430-1350		
f. S-O stretching vibrations; sulfonates		8.44-8.59	(s)	1155-1165	
and 7.50-7.68		(s)	1370-1340		
g. S-O stretching vibrations; sulfonamides	8.44-8.77	(s)	1180-1140		
and 7.41-7.59	(s)	1350-1300			
h. S-O stretching vibrations; sulfonic acids	8.27-8.20	(s)	1210-1150		
and 9.43-9.71	(s)	1060-1030			
and 13.5	(s)	~650			

f. Abbreviations: s = strong, m = medium, w = weak, v = variable, b = broad, sh = sharp.
- = approximately

Proton Chemical Shifts (Values are given on the officially approved δ scale; $\tau = 10.00 - \delta$)

Substituent Group	Methyl Protons	Methylene Protons	Methine Proton
HC-C-CH ₃	0.95	1.20	1.55
HC-C-NR ₂	1.05	1.45	1.70
HC-C-C=C	1.00	1.35	1.70
HC-C-C=O	1.05	1.55	1.95
HC-C-NR ₂ Ar	1.10	1.50	1.80
HC-C-NH(C=O)R	1.10	1.50	1.90
HC-C-(C=O)NR ₂	1.10	1.50	1.80
HC-C-(C=O)Ar	1.15	1.55	1.90
HC-C-(C=O)OR	1.15	1.70	1.90
HC-C-Ar	1.15	1.55	1.90
HC-C-OH (and OR)	1.20	1.50	1.80
HC-C-C=CR	1.20	1.50	1.75
HC-C-C=N	1.25	1.65	1.80
HC-C-SR	1.25	1.60	2.00
HC-C-OAr	1.30	1.55	1.90
HC-C-O(C=O)R	1.30	1.60	2.00
HC-C-SH	1.30	1.60	1.80
HC-C-(S=O)R and -SO ₂ R	1.35	1.60	1.65
HC-C-NR ₂	1.40	1.75	2.05
HC-C-O(C=O)CF ₃	1.40	1.65	1.95
HC-C-Cl	1.55	1.80	1.85
HC-C-O(C=O)Ar	1.65	1.75	1.90
HC-C-Br	1.80	1.85	1.90
HC-CH ₃	0.90	1.30	1.50
HC-C=C	1.60	2.05	2.80
HC-C=C	1.70	2.20	2.50
HC-(C=O)OR (and NR ₂)	2.00	2.35	3.00
HC-SR	2.05	2.55	2.55
HC-O-O	2.10	2.30	2.55
HC-(C=O)R	2.10	2.35	2.65
HC-C=N	2.15	2.45	2.90
HC-CHO	2.20	2.40	2.85
HC-Ar (and NR ₂)	2.25	2.45	3.40
HC-SR	2.35	2.70	3.60
HC-(C=O)Ar	2.40	2.70	3.60
HC-SAr	2.40	3.10	4.10
HC-NR ₂ Ar	2.60	3.05	3.60
HC-SO ₂ R and -(SO) ₂ R	2.70	3.40	4.10
HC-Br	2.70	3.15	3.85
HC-NR ₂	2.95	3.35	4.05
HC-NH(C=O)R	3.05	3.45	3.60
HC-Cl	3.20	3.40	4.05
HC-OH and -OR	3.50	3.75	4.95
HC-O(C=O)R	3.65	4.10	4.60
HC-OAr	3.80	4.00	5.05
HC-O(C=O)Ar	3.80	4.20	

TURN OVER

Substituent Group	Methyl Protons	Methylene Protons	Methine Proton
HC-F	4.25	4.50	4.80
HC-NO ₂	4.30	4.55	4.60
Cyclopropane	-0.20	2.45	0.40
Cyclobutane	2.45	1.65	
Cyclopentane	1.50	1.25	
Cyclohexane	1.50	1.25	
Cycloheptane	1.50	1.25	

9

Substituent Group	Proton Shift	Substituent Group	Proton Shift
HC=CH	2.35	HO-C=O	10-12
HC=CAr	2.90	HO-SO ₂	11-12
HC=C-C=C	2.75	HO-Ar	4.5-6.5
HAr	7.20	HO-R	0.5-4.5
HCO-O	8.1	HS-Ar	2.8-3.6
HCO-R	9.4-10.0	HS-R	1-2
HCO-Ar	9.7-10.5	HN-Ar	3-5
HQ-N=C(O)Ar	9-12	HN-R	0.5-5

¹³C Chemical Shifts (Values given on the δ scale, relative to TMS.)

Substituent Group	Primary Carbon	Secondary Carbon	Tertiary Carbon	Quaternary Carbon
Alkenes	-20 to 30	25 to 45	30 to 60	35 to 70
C-C=C	40 to 60	49 to 70	60 to 75	70 to 85
C-O	20 to 45	40 to 60	50 to 70	55 to 75
C-N	10 to 30	25 to 45	40 to 55	55 to 70
C-S	10 to 30	10 to 45	30 to 65	35 to 75
C-Halide	-17 to 35	(1) (C)	(1) (C)	(1) (C)
Alkyne	70 to 100	130 to 150	150 to 160	150 to 160
Alkenes	110 to 150	130 to 150	150 to 165	150 to 165
Aromatics	110 to 135	130 to 150	150 to 170	150 to 170
Cy-substituted	125 to 145	130 to 150	150 to 170	150 to 170
Heteroaromatics	115 to 140	130 to 150	150 to 170	150 to 170
C=O	135 to 155	130 to 150	150 to 170	150 to 170
Cyanates	105 to 120	130 to 150	150 to 170	150 to 170
Isocyanates	115 to 135	130 to 150	150 to 170	150 to 170
Thiocyanates	110 to 130	130 to 150	150 to 170	150 to 170
Isothiocyanates	120 to 140	130 to 150	150 to 170	150 to 170
Cyanides	110 to 130	130 to 150	150 to 170	150 to 170

SPIN-SPIN COUPLING CONSTANTS

Type	J, cps	Type	J, cps
H_2^+ CH_2^+ 	280	$C-CH=CH-C$ 	9-13
$C-H$ 	12-4	$H-C\equiv C-H$ 	9.1
$CH-CH$ 	12-15	$CH-C\equiv C-H$ 	2-3
$-C-(C)-C-$ 	2.9	$CH-C$ 	1-3
CH_2-CH_2-X 	~0	$C=C$ 	6-8
$CH-N$ 	6.5-7.5		<i>o</i> -6-9 <i>m</i> -1-3 <i>p</i> -0-1
CH_2 	5.5-7.0		<i>ortho</i> 1.6-2.0 <i>meta</i> 0.6-1.0 <i>para</i> 1.3-1.8 <i>ortho</i> 3.2-3.8
$H-C-C-H$ 	<i>aa</i> 5-10 <i>ac</i> 2-3 <i>cc</i> 2-4		<i>ortho</i> 2.0-2.6 <i>meta</i> 1.5-2.2 <i>para</i> 1.8-2.3 <i>ortho</i> 2.8-4.0
$C=C$ 	0.5-3		<i>ortho</i> 4.6-5.8 <i>meta</i> 1.0-1.8 <i>para</i> 2.1-3.3 <i>ortho</i> 5.0-4.5
$C=C$ 	7-12		<i>ortho</i> 4.9-5.7 <i>meta</i> 1.6-2.6 <i>ortho</i> 0.7-1.1 <i>meta</i> 0.2-0.5 <i>ortho</i> 7.2-8.5 <i>ortho</i> 1.4-1.9
$C=C$ 	13-18		
$C=C$ 	4-10		
$C=C$ 	0.5-2.5		
$C=C$ 	~0		

TURN OVER

Conjugated dienes & Trienes, Solvent: Ethanol

Parent value for Butadiene system or acyclic conjugated diene	217 nm
Acyclic triene	245 nm
Homoannular conjugated diene	253 nm
Heteroannular conjugated diene	215 nm
Increment for each substituents	
Alkyl substituents or ring residue	5 nm
Exocyclic double bond	5 nm
Double bond extending conjugation	30 nm
Auxochrome	
-OR	+6 nm
-SR	+30 nm
-Cl, -Br	+5 nm
NR ₂	+60 nm
-OCOCH ₃	0

Woodward Fieser rules for $\alpha\beta$ -unsaturated carbonyl compounds:-

a) Parent values			
$\alpha\beta$ -unsaturated acyclic or six membered ketone	215 nm		
$\alpha\beta$ -unsaturated five membered ring ketone	202 nm		
$\alpha\beta$ -unsaturated aldehyde	207 nm		
b) Increments			
i) Each alkyl substituents or ring residue			
At α position	10 nm		
At β position	12 nm		
At gamma and higher position	18 nm		
ii) Each exocyclic double bond	5 nm		
iii) Double bond extending conjugation	30 nm		
iv) Homoannular conjugated diene	39 nm		
Auxochromes	Positions		
	α	β	gamma
-OH	35	30	50
-OR	35	30	17
-SR	-	85	-
-OCOCH ₃	6	6	6
-Cl	15	12	-
-Br	25	30	-
-NR ₂	-	95	-